

TEN MOST IMPORTANT PUBLICATIONS

Schäfer I.B., Yamashita M., Schuller J.M., Schüssler S., Reichelt P., Strauss M. and **Conti** E. (2019). Molecular basis for poly(A) RNP architecture and recognition by the Pan2-Pan3 deadenylase. *Cell* 177, 1619-1631.

Schuller J. M., Falk S., Fromm L., Hurt E. and **Conti** E. (2018). Structure of the nuclear exosome captured on a maturing pre-ribosome. *Science* 360, 219-222.

Gerlach P, Schuller JM, Bonneau F, Basquin J, Reichelt P, Falk S and **Conti** E. (2018) Distinct and evolutionary conserved structural features of the human nuclear exosome complex. *Elife* 7:e38686

Schmidt C., Kowalinski E., et al. **Conti*** E., Becker* T and Beckmann* R. (2017). The cryo-EM structure of a ribosome – Ski2-Ski3-Ski8 complex. *Science*, 354, 1431-1433.

Makino D.L., Schuch B., Stegmann E., Baumgärtner M., Basquin C. and **Conti** E. (2015). RNA degradation paths in a 12-subunit nuclear exosome complex. *Nature* 524, 54-58.

Halbach F., Reichelt P. Rode M. and **Conti** E. (2013). The yeast Ski complex: crystal structure and substrate channeling to the exosome. *Cell* 154, 814-826

Makino D.L., Baumgärtner M. and **Conti** E. (2013). Crystal structure of an RNA-bound 11-subunit eukaryotic exosome complex. *Nature* 495, 70-75.

Bonneau F., Basquin J., Ebert J., Lorentzen E. and **Conti** E. (2009). The yeast exosome functions as a macromolecular cage to channel RNA substrates for degradation. *Cell* 139, 547-559.

Cook A., Fukuhara N., Jinek M. and **Conti** E. (2009). Structures of the tRNA export factor in the nuclear and cytosolic states. *Nature* 461, 60-65.

Bono F., Ebert J., Lorentzen E., and **Conti** E. (2006). The crystal structure of the exon junction complex reveals how it maintains a stable grip on mRNA. *Cell* 126, 713-725.

COMPLETE LIST OF PUBLICATIONS

Schuller S.K., Schuller J.M., Prabu J.R., Baumgärtner M., Bonneau F., Basquin J., **Conti** E. (2020). Structural insights into the nucleic acid remodeling mechanisms of the yeast THO-Sub2 complex. *eLife* 9:e61467.

Lingaraju M., Schuller J.M., Falk S., Gerlach P., Bonneau F., Basquin J., Benda C., **Conti** E. (2020). To Process or to Decay: A Mechanistic View of the Nuclear RNA Exosome. *Cold Spring Harb Symp Quant Biol* 2019, 84:155-163.

Langer L.M., Gat Y., Bonneau F., **Conti** E. (2020). Structure of substrate-bound SMG1-8-9 kinase complex reveals molecular basis for phosphorylation specificity. *Elife*, 9 e57127.

Busetto V., Barbosa I., Basquin J., Marquenet É., Hocq R., Hennion M., Paternina J.A., Namane A., **Conti** E., Bensaude O., Le Hir H. (2020). Structural and functional insights into CWC27/CWC22 heterodimer linking the exon junction complex to spliceosomes. *Nucleic Acids Res.* 48, 5670-5683.

Keidel A., **Conti** E. and Falk S. (2020). Purification and Reconstitution of the *S. cerevisiae* TRAMP and Ski Complexes for Biochemical and Structural Studies. *Methods Mol Biol.* 2062, 491-513.

Gat Y., Schuller J.M., Lingaraju M., Weyher E., Bonneau F., Strauss M., Murray P.J. and **Conti** E. (2019). InsP6 binding to PIKK kinases revealed by the cryo-EM structure of an SMG1-SMG8-SMG9 complex. *Nat Struct Mol Biol.* 26, 1089-1093.

Lingaraju M., Johnsen D., Schlundt A., Langer L.M., Basquin J., Sattler M., Jensen T.H., Falk S. and **Conti** E. (2019) The MTR4 helicase recruits nuclear adaptors of the human exosome using distinct arch-interacting motifs. *Nature Communications* 10, 3393.

Schäfer I.B., Yamashita M., Schuller J.M., Schüssler S., Reichelt P., Strauss M. and **Conti** E. (2019). Molecular basis for poly(A) RNP architecture and recognition by the Pan2-Pan3 deadenylase. *Cell* 177, 1619-1631.

Gerlach P, Schuller JM, Bonneau F, Basquin J, Reichelt P, Falk S and **Conti** E. (2018) Distinct and evolutionary conserved structural features of the human nuclear exosome complex. *Elife* 7:e38686

Schuller J. M., Falk S., Fromm L., Hurt E. and **Conti** E. (2018). Structure of the nuclear exosome captured on a maturing pre-ribosome. *Science* 360, 219-222.

Fromm L., Falk S., Flemming D., Schuller J.M., Thoms M., **Conti** E. and Hurt E. (2017). Reconstitution of the complete pathway of ITS2 processing at the pre-ribosome. *Nature Commun*, 8, 1787.

Falk S., Bonneau F., Ebert J., Kögel A. and **Conti** E. (2017). Mpp6 incorporation into the nuclear exosome contributes to RNA channeling through the Mtr4 helicase. *Cell Rep* 20, 2279-2286.

Li L., Lingaraju M., Basquin C. Basquin J. and **Conti** E. (2017). Structure of a SMG8-SMG9 complex identifies a G-domain heterodimer in the NMD effector proteins. *RNA* 23, 1028-1034.

Leonaitė B., Han Z., Basquin J., Bonneau F., Libri D., Porrua O. and **Conti** E. (2017). Sen1 has unique structural features grafted on the architecture of the Upf1-like helicase family. *EMBO J.* 36, 1590-1604.

Falk S., Tants J.-N., Thoms M., Hurt E., Sattler M. and **Conti** E. (2017). Structural insights into the interaction of the nuclear exosome helicase Mtr4 with the pre-ribosomal protein Nop53. *RNA* 23, 1780-1787.

Schmidt C., Kowalinski E., Shanmuganathan V., Defenouillère Q., Braunger K., Heuer A., Pech M., Namane A., Berninghausen O., Fromont-Racine M., Jacquier A., **Conti*** E., Becker* T and Beckmann* R. (2017). The cryo-EM structure of a ribosome – Ski2-Ski3-Ski8 complex. *Science*, 354, 1431-1433. (* co-corresponding authors)

Falk S., Finogenova K., Melko M., Benda C., Lykke-Andersen S., Heick Jensen T. and **Conti** E. (2016). Crystal structure of a core RRM-protein interaction in the human NEXT complex reveals a physical connection to splicing factors. *Nature Comm.*, 7, 13573.

Kowalinski E., Koegel A., Ebert J., Reichelt P., Stegmann E., Habermann B. and **Conti** E. (2016). Structure of a cytoplasmic 11-subunit RNA exosome complex. *Mol. Cell* 63, 125-134.

Nakel K., Bonneau F., Basquin C., Habermann B., Eckmann R.E. and **Conti** E. (2016). Structural basis for the antagonistic roles of RNP-8 and GLD-3 in GLD-2 poly(A)-polymerase activity. *RNA* 22, 1139-1145.

Vukovic L., Chipot C., Makino D.L., **Conti** E. and Schulten K. (2016). Molecular mechanism of processive 3' to 5' RNA translocation in the active subunit of the RNA exosome complex. *J. Am. Chem Soc.* 138, 4069-4078.

Prabu J.R., Müller M., Thomae A.W., Schüssler S., Bonneau F., Becker P.B. and **Conti** E. (2015). Structure of the RNA helicase MLE reveals the molecular mechanisms for uridine specificity and RNA-ATP coupling. *Mol. Cell* 60, 487-499.

Ozgur S., Basquin J., Kamenska A., Filipowicz W., Standart N. and **Conti** E. (2015). Structure of a human 4E-T - DDX6 - CNOT1 complex reveals the different interplay of DDX6-binding proteins with the CCR4-NOT complex. *Cell Rep.* 13, 703-711.

Makino D.L., Schuch B., Stegmann E., Baumgärtner M., Basquin C. and **Conti** E. (2015). RNA degradation paths in a 12-subunit nuclear exosome complex. *Nature* 524, 54-58.

Nakel K., Bonneau F., Eckmann C.R. and **Conti** E. (2015). Structural basis for the activation of the *C. elegans* noncanonical cytoplasmic polymerase GLD-2 by GLD-3. *Proc. Natl. Acad. USA* 112, 86414-86419.

Kowalinski E., Schuller A., Green R. and **Conti** E. (2015). *S. cerevisiae* Ski7 is a GTP-binding protein adopting the characteristic conformation of active translational GTPases. *Structure* 23, 1336-1343.

Bhaskar V., Basquin J. and **Conti** E. (2015). Architecture of the ubiquitylation module of the yeast Ccr4-Not complex. *Structure* 23, 921-928.

Ozgur S., Buchwald, G., Falk S., Chakrabarti S., Prabu J.R. and **Conti** E. (2015). The conformational plasticity of eukaryotic RNA-dependent ATPases. *FEBS J.* 282, 850-863.

Schuch B., Feigenbutz M., Makino D.L., Falk S., Basquin C., Mitchell P. and **Conti** E.^[SEP] (2014) The exosome-binding factors Rrp6 and Rrp47 form a composite surface for recruiting the Mtr4

helicase.^[L]_[SEP] *EMBO J.* 33, 2829-2846.

Benda C., Ebert J., Scheltema R.A., Schiller H.B., Baumgärtner M., Bonneau F., Mann M. and **Conti** E. (2014). ^[L]_[SEP]Structural Model of a CRISPR RNA-Silencing Complex Reveals the RNA-Target Cleavage Activity in Cmr4.^[L]_[SEP] *Mol. Cell* 56, 43-54.

Falk S., Weir J., Hentschel J., Reichelt P., Bonneau F. and **Conti** E. (2014). The molecular architecture of the TRAMP complex reveals the organization and interplay of its two catalytic activities. *Mol. Cell* 55, 856-867.

Chakrabarti S., Bonneau F., Schüssler S., Eppinger E. and **Conti** E. (2014) ^[L]_[SEP]Phospho-dependent and phospho-independent interactions of the helicase UPF1 with the NMD factors SMG5-SMG7 and SMG6.^[L]_[SEP] *Nucleic Acids Res.* 42, 9447-9460.

Schäfer I.B., Rode M., Bonneau F., Schüssler S. and **Conti** E. (2014). The structure of the Pan2-Pan3 core complex reveals cross-talk between deadenylase and pseudokinase.^[L]_[SEP] *Nature Struct. Mol. Biol.* 21, 591-598.

Mathys H.*^[L], Basquin J.*^[L], Ozgur S., Czarnocki-Cieciura M., Bonneau F., Aartse A., Dziembowski A., Nowotny M., **Conti** E.*^[L], Filipowicz W.*^[L]. (2014). Structural and Biochemical Insights to the Role of the CCR4-NOT Complex and DDX6 ATPase in MicroRNA Repression.^[L]_[SEP] *Mol. Cell* 54, 751-765. (* co-corresponding authors)

Buchwald G. Schüssler S, Basquin. C., Le Hir H. and **Conti** E. (2013). Crystal structure of the human eIF4AIII-CWC22 complex shows how a DEAD-box protein is inhibited by a MIF4G domain. *Proc. Natl. Acad. USA* 110, E4611-E4618.

Bhaskar V., Roudko V., Basquin J., Sharma K., Urlaub H., Seraphin B. and **Conti** E. (2013). Structure and RNA-binding properties of the Not1–Not2–Not5 module of the yeast Ccr4–Not complex. *Nature Struct. Mol. Biol.* 20, 1281-1288.

Sharif H. and **Conti** E. (2013). Architecture of the Lsm1-7-Pat1 Complex: a conserved assembly in eukaryotic mRNA turnover. *Cell Rep.* 5, 283-291.

Makino D.L., Halbach F. and **Conti** E. (2013). The exosome and proteasome: common principles of degradation control. *Nature Rev. Mol. Cell. Biol.* 14, 650-660. (review)

Halbach F., Reichelt P., Rode M. and **Conti** E. (2013). The yeast Ski complex: crystal structure and substrate channeling to the exosome. *Cell* 154, 814-826.

Makino D.L. and **Conti** E. (2013) Structure determination of an 11-subunit exosome in complex with RNA by Molecular Replacement. *Acta Cryst. D*69, 2226-2235.

Sharif H., Ozgur S., Sharma K., Basquin C., Urlaub H. and **Conti** E. (2013). Structural analysis of the yeast Dhh1-Pat1 complex reveals how Dhh1 engages Pat1, Edc3 and RNA in mutually exclusive interactions. *Nucleic Acids Res.* 41, 8377-8390.

Von Moeller H., Lerner R., Ricciardi A., Basquin C., Marzluff W.F. and **Conti** E. (2013). Structural and biochemical studies of SLIP1-SLBP identifies DBP5 and eIF3g as SLIP1-binding proteins. *Nucleic Acids Res.* 41, 7960-7971.

Makino D.L., Baumgärtner M. and **Conti** E. (2013). Crystal structure of an RNA-bound 11-subunit eukaryotic exosome complex. *Nature* 495, 70-75.

Basquin J., Roudko V.V., Rode M., Basquin C., Seraphin B. and **Conti** E. (2012). Architecture of the nuclease module of the Ccr4-Not complex: the Not1-Caf1-Ccr4 interaction. *Mol. Cell* 48, 1-12.

Melero R., Buchwald G., Castano R., Raabe M., Gil D., Lazaro M., Urlaub, H., **Conti** E. and Llorca O. (2012). The cryo-EM structure of the UPF-EJC complex shows UPF1 is poised toward the RNA 3' end. *Nature Struct. Mol. Biol.* 19, 498-505.

Jeyaprakash A.A., Santamaria A., Jayachandran U., Chan Y.W., Benda C., Nigg E. and **Conti** E. (2012). Structural and functional organization of the Ska complex, a key component of the kinetochore-microtubule interface. *Mol. Cell* 46, 1-13.

Murachelli A.G., Ebert J., Basquin C., Le Hir H. and **Conti** E. (2012). The structure of the ASAP core complex reveals the existence of a Pinin-containing PSAP complex. *Nature Struct. Mol. Biol.* 19, 378-386.

Halbach F., Rode M. and **Conti** E. (2012). The crystal structure of *S. cerevisiae* Ski2, a DExH helicase associated with the cytoplasmic functions of the exosome. *RNA* 18, 124-134.

Jeyaprakash A.A., Basquin C., Jayachandran U. and **Conti** E. (2011). Structural basis for the recognition of phosphorylated histone H3 by the Survivin subunit of the Chromosomal Passenger Complex. *Structure* 19, 1-10.

Chakrabarti S., Jayachandran U., Bonneau F., Fiorini F., Basquin C., Domcke S., Le Hir H. and **Conti** E. (2011). Molecular mechanisms for the RNA-dependent ATPase activity of Upf1 and its regulation by Upf2. *Mol. Cell* 41, 693-703.

Malet H., Topf M., Clare D.K., Ebert J., Bonneau F., Basquin J., Drazkowska K., Tomecki R., Dziembowski A., **Conti** E., Sabil H.R. and Lorentzen E. (2010). RNA channeling by the eukaryotic exosome. *EMBO Rep.* 11, 936-942.

Nakel K., Hartung S.A., Bonneau F., Eckmann C.R. and **Conti** E. (2010). Four KH domains of the *C. elegans* Bicaudal-C ortholog GLD-3 form a globular structural platform. *RNA* 16, 2058-2067.

Kashima I., Jonas S., Jayachandran U., Buchwald G., **Conti** E., Lupas A.N. and Izaurralde E. (2010). SMG6 interacts with the exon junction complex via two conserved EJC-binding motifs (EBMs) required for nonsense-mediated mRNA decay. *Genes and Dev.* 24, 2440-2450.

Weir J.R., Bonneau F., Hentschel J. and **Conti** E. (2010). Structural analysis reveals the characteristic features of Mtr4, a DExH helicase involved in nuclear RNA processing and surveillance. *Proc. Natl. Acad. USA* 107, 12139-12144.

Buchwald G., Ebert J., Basquin C., Sauliere J., Jayachandran U., Bono F., Le Hir H. and **Conti** E. (2010). Insights into the recruitment of the NMD machinery from the crystal structure of a core EJC-Upf3b complex. *Proc. Natl. Acad. USA* 107, 10050-10055.

Cook A.G. and **Conti** E. (2010). Nuclear export complexes in the frame. *Current Opinion Struct. Biol.* 20, 247-252. (review)

Bono F., Cook A., Grünwald M., Ebert J. and **Conti** E. (2010). Nuclear import mechanism of the EJC component Mago-Y14 revealed by structural studies of Importin13. *Mol. Cell* 37, 211-222.

Bonneau F., Basquin J., Ebert J., Lorentzen E. and **Conti** E. (2009). The yeast exosome functions as a macromolecular cage to channel RNA substrates for degradation. *Cell* 139, 547-559.

Cook A., Fukuhara N., Jinek M. and **Conti** E. (2009). Structures of the tRNA export factor in the nuclear and cytosolic states. *Nature* 461, 60-65.

Von Moeller H., Basquin C. and **Conti** E. (2009). The mRNA export protein Dbp5 binds RNA and the cytoplasmic nucleoporin Nup214 in a mutually exclusive manner. *Nature Struct. Mol. Biol.* 16, 247-254.

Lorentzen E., Basquin J. and **Conti** E. (2008). Structural organization of the RNA degrading exosome. *Current Opinion Struct. Biol.* 18, 709-713. (review)

Jinek M., Eulalio A., Lingel A., Helms S., **Conti** E. and Izaurrealde, E. (2008). The C-terminal region of Ge-1 presents conserved structural features required for P-body localization. *RNA* 14, 1991-1998.

Lorentzen E., Basquin J., Tomecki R., Dziembowski A. and **Conti** E. (2008). Structure of the active subunit of the yeast exosome core, Rrp44: diverse modes of substrate recruitment in the RNase II nuclease family. *Mol. Cell* 29, 717-728.

Jeyaprakash A.A., Klein U.R., Lindner D., Ebert J., Nigg E.A. and **Conti** E. (2007). Structure of a Survivin-Borealin-INCENP core complex reveals how Chromosomal Passengers travel together. *Cell* 131, 271-285.

Cook A., Bono F., Jinek M. and **Conti** E. (2007). Structural biology of nucleocytoplasmic transport. *Annual Review Biochem.* 76, 647-671. (review)

Lorentzen E., Dziembowski A., Lindner D., Seraphin B. and **Conti** E. (2007). RNA channelling by the archaeal exosome. *EMBO Rep.* 8, 470-476.

Dziembowski A., Lorentzen E., **Conti** E. and Seraphin B. (2007). A single subunit, Dis3, is essentially responsible for the yeast exosome activity. *Nature Struct. Mol. Biol.* 14, 15-22.

Oddone A., Lorentzen E., Basquin J., Gasch A., Rybin V., **Conti** E. and Sattler S. (2007). Structural and biochemical characterization of the yeast exosome component Rrp40. *EMBO Rep.* 8, 63-69.

Glavan F., Behm-Ansmant I., Izaurrealde E. and **Conti** E. (2006). Structures of the PIN domains of SMG6 and SMG5 reveal a nuclease within the mRNA surveillance complex. *EMBO J.* 25, 5117-5125.

Bono F., Ebert J., Lorentzen E. and **Conti** E. (2006). The crystal structure of the exon junction complex reveals how it maintains a stable grip on mRNA. *Cell* 126, 713-725.

Jinek M., Chen Y.W., Clausen H., Cohen S.M. and **Conti** E. (2006). Structural insights into the Notch-modifying glycosyltransferase Fringe. *Nature Struct Mol Biol.* 13, 945-946.

Jinek M and **Conti** E. (2006). Eukaryotic expression, purification, crystallization and preliminary X-ray analysis of murine Manic Fringe. *Acta Cryst. Sect F* 62, 774-777.

Lorentzen E. and **Conti** E. (2006). The exosome and the proteasome: nano-compartments for degradation. *Cell* 125, 651-654. (review)

Conti E., Muller C.W. and Stewart M. (2006). Karyopherin flexibility in nucleocytoplasmic transport. *Current Opinion Struct Biol.* 16, 237-244. (review)

Lorentzen E. and **Conti** E. (2005). Structural basis of 3'-end RNA recognition and exoribonucleolytic cleavage by an exosome RNase PH core. *Mol. Cell* 20 473-481.

Lorentzen E., Walter P., Fribourg S., Evguenieva-Hackenberg E., Klug G. and **Conti** E. (2005). The archaeal exosome core is a hexameric ring with three catalytic subunits. *Nature Struct. Mol. Biol.* 12, 575-581.

Conti E. and Izaurralde E. (2005). Nonsense-mediated mRNA decay: molecular insights and mechanistic variations across species. *Current Opinion Cell Biol.* 17, 316-325 (review).

Cook A., Fernandez E., Lindner D., Ebert J., Schlenstedt G. and **Conti** E. (2005). The structure of the nuclear export receptor Cse1 in its cytosolic state reveals a closed conformation incompatible with cargo binding. *Mol. Cell* 18, 355-367.

Fukuhara N., Ebert J., Unterholzner L., Lindner D., Izaurralde E. and **Conti**, E. (2005). SMG7 is a 14-3-3-like adaptor in the nonsense-mediated mRNA decay pathway. *Mol. Cell* 17, 537-547.

Jinek M., Rehwinkel J., Lazarus B.D., Izaurralde E., Hanover J.A. and **Conti** E. (2004). The superhelical TPR-repeat domain of O-linked GlcNAc transferase exhibits structural similarities to importin α . *Nature Struct. Mol. Biol.* 11, 1001-1007.

Bono F., Ebert E., Unterholzner L., Guttler T., Izaurrealde E. and **Conti**, E. (2004). Molecular insights into the interaction of PYM with the Mago-Y14 core of the exon junction complex. *EMBO Rep.* 5, 304-310.

Bayliss R., Sardon T., Ebert J., Lindner D., Vernos I. and **Conti** E. (2004). Determinants for Aurora-A activation and Aurora-B discrimination by TPX2. *Cell Cycle* 3, 404-407.

Fukuhara N., Fernandez E., Ebert J., **Conti** E. and Svergun D. (2004). Conformational variability of nucleo-cytoplasmic transport factors. *J. Biol. Chem.* 279, 2176-2181.

Bayliss R., Sardon T., Vernos I. and **Conti** E. (2003). Structural basis of Aurora-A activation by TPX2 at the mitotic spindle. *Mol. Cell* 12, 851-862.

Fribourg S. and **Conti** E. (2003). Structural similarity in the absence of sequence homology of the mRNA export factors Mtr2 and p15. *EMBO Rep.* 4, 699-703.

Fribourg S., Gatfield D., Izaurrealde E. and **Conti** E. (2003). A novel mode of RBD-protein recognition in the Y14-mago complex. *Nature Struct. Biol.* 10, 433-439.

Fribourg S., Braun I.C., Izaurrealde E. and **Conti** E. (2001). Structural basis for the recognition of a nucleoporin FG repeat by the NTF2-like domain of the TAP/p15 mRNA nuclear export factor. *Mol. Cell* 8, 645-56.

Braun I.C., Herold A., Rode M., **Conti** E. and Izaurrealde E. (2001). Overexpression of TAP/p15 heterodimers bypasses nuclear retention and stimulates nuclear mRNA export. *J. Biol. Chem.* 276, 20536-20543.

Conti E. and Izaurrealde E. (2001). Nucleocytoplasmic transport enters the atomic age. *Current Opinion Cell Biol.* 13, 310-319 (review).

Liker E., Fernandez E., Izaurrealde E. and **Conti** E. (2000). The structure of the RNA export factor TAP reveals a non-canonical RNP in cis with a LRR domain. *EMBO J.* 19, 5587-5598.

Conti E. and Kuriyan J. (2000). Crystallographic analysis of the specific yet versatile recognition of distinct nuclear localization signals by karyopherin α . *Structure* 8, 329-338.

Mattaj I.W. and **Conti** E. (1999). Cell biology. Snail mail to the nucleus. *Nature* 399, 208-210. (review).

Conti E., Uy M., Leighton L., Blobel G. and Kuriyan J. (1998). Crystallographic analysis of the recognition of a nuclear localization signal by the nuclear import factor karyopherin- α . *Cell* 94, 193-204.

Conti E., Stachelhaus T., Marahiel M. and Brick P. (1997). Structural basis for the activation of phenylalanine in the non-ribosomal synthesis of gramicidin S. *EMBO J.* 16, 4174-4183.

Conti E., Rivetti C., Wonacott A. and Brick P. (1998). X-ray and spectrophotometric studies of the binding of proflavin to the S1 specificity pocket of human α -thrombin. *FEBS Letters* 425, 229-233.

Conti E., Franks N.P. and Brick P. (1996). Crystal structure of firefly luciferase throws light on a superfamily of adenylate-forming enzymes. *Structure* 4, 287-298.

Conti E., Moser C., Rizzi M., et al. and Bolognesi M. (1993). X-ray crystal structure of ferric *Aplysia limacina* myoglobin in different liganded states. *J. Mol. Biol.* 233, 498-508.